

1. Гомотопійна категорія категорії модулів.  
Навколо похідних категорій

Володимир Любашенко

4 лютого 2021

# Категорії модулів $\mathcal{C} = \mathbb{k}\text{-mod}$ . Коланцюгові комплекси

Let  $\mathcal{C}$  be an additive category.

**Definition 1.3.1.** A complex  $X$  in  $\mathcal{C}$  consists of the data  $\{X^n, d_X^n\}_{n \in \mathbb{Z}}$ , such that for any  $n \in \mathbb{Z}$ :

$$(1.3.1) \quad X^n \in \text{Ob}(\mathcal{C}), \quad d_X^n \in \text{Hom}_{\mathcal{C}}(X^n, X^{n+1}) \quad \text{and} \quad d_X^{n+1} \circ d_X^n = 0 .$$

A morphism  $f$  from a complex  $X$  to a complex  $Y$  is a sequence  $\{f^n\}_{n \in \mathbb{Z}}$  of morphisms  $f^n: X^n \rightarrow Y^n$ , such that for any  $n$ :

$$(1.3.2) \quad d_Y^n \circ f^n = f^{n+1} \circ d_X^n .$$

We denote by  $\mathbf{C}(\mathcal{C})$  the category of complexes of  $\mathcal{C}$  thus obtained.

One often writes a complex as a sequence:

$$\cdots \longrightarrow X^{n-1} \xrightarrow{d_X^{n-1}} X^n \xrightarrow{d_X^n} X^{n+1} \longrightarrow \cdots .$$

The family  $d_X = \{d_X^n\}_n$  is called the **differential** of the complex  $X$ . A complex  $X$  is said to be **bounded** (resp. **bounded below**, resp. **bounded above**) if  $X^n = 0$  for  $|n| \gg 0$  (resp.  $n \ll 0$ , resp.  $n \gg 0$ ). The full subcategory of  $\mathbf{C}(\mathcal{C})$  consisting of bounded complexes (resp. complexes bounded below, resp. complexes bounded above), is denoted  $\mathbf{C}^b(\mathcal{C})$  (resp.  $\mathbf{C}^+(\mathcal{C})$ , resp.  $\mathbf{C}^-(\mathcal{C})$ ).

We identify  $\mathcal{C}$  with the full subcategory of  $\mathbf{C}(\mathcal{C})$  consisting of complexes  $X$  such that  $X^n = 0$  for  $n \neq 0$ .

## Функтор зсуву

**Definition 1.3.2.** Let  $k$  be an integer, and let  $X \in \text{Ob}(\mathbf{C}(\mathcal{C}))$ . One defines a new complex  $X[k]$  by setting:

$$(1.3.3) \quad \begin{cases} X[k]^n = X^{n+k} , \\ d_{X[k]}^n = (-1)^k d_X^{n+k} . \end{cases}$$

For a morphism  $f : X \rightarrow Y$  in  $\mathbf{C}(\mathcal{C})$ , one defines  $f[k] : X[k] \rightarrow Y[k]$  by setting:

$$(1.3.4) \quad f[k]^n = f^{n+k} .$$

The functor  $[k]$  from  $\mathbf{C}(\mathcal{C})$  to  $\mathbf{C}(\mathcal{C})$  is called the **shift functor** of degree  $k$ .

For any graded  $\mathbb{k}$ -module  $M$  and an integer  $a$  denote by  $M[a]$  the same module with the grading shifted by  $a$ :  $M[a]^k = M^{a+k}$ .

Denote by  $\sigma^a : M \rightarrow M[a]$ ,  $M^k \ni x \mapsto x \in M[a]^{k-a}$  the “identity map” of degree  $\text{deg } \sigma^a = -a$ . Write elements of  $M[a]$  as  $m\sigma^a$ .

Typically, a map is written on the right of its argument. When  $f : V \rightarrow X$  is a homogeneous map of certain degree, the map  $f[a] : V[a] \rightarrow X[a]$  is defined as

$$f[a] = (-1)^{a \text{deg } f} \sigma^{-a} f \sigma^a = (-1)^{af} \sigma^{-a} f \sigma^a .$$

Нехай  $K$ -комутативне кільце  $\in \mathbb{Z}$ .

Тензорний добуток  $K$ -модулів  $(M, N) \mapsto M \otimes_K N$  перетворює  $K$ -мод на симетричну моноїдальну категорію.

Вона замкнена зліва і справа, тобто  $\exists K$ -модулі  $\underline{K\text{-mod}}^2(M, N)$  і  $\underline{K\text{-mod}}^0(M, N)$  разом з паруваними  $ev^2: M \otimes \underline{K\text{-mod}}^2(M, N) \rightarrow N$

$$ev^2: \underline{K\text{-mod}}^0(M, N) \otimes M \rightarrow N \quad \text{Тож}$$

$\forall L \in K\text{-mod} \quad \forall \varphi: M \otimes L \rightarrow N \exists! \xi: L \rightarrow \underline{K\text{-mod}}^2(M, N) :$

$$\begin{array}{ccc} M \otimes L & \xrightarrow{\varphi} & N \\ \text{id} \otimes \downarrow & \searrow & \\ M \otimes \underline{K\text{-mod}}^2(M, N) & \xrightarrow{ev^2} & N \end{array}$$

$\forall L \in K\text{-mod} \quad \forall \chi: L \otimes M \rightarrow N \exists! \xi: L \rightarrow \underline{K\text{-mod}}^0(M, N) :$

$$\begin{array}{ccc} L \otimes M & \xrightarrow{\chi} & N \\ \text{id} \otimes \downarrow & \searrow & \\ \underline{K\text{-mod}}^0(M, N) \otimes M & \xrightarrow{ev^2} & N \end{array}$$

$$M \otimes M$$

$$\underline{K\text{-mod}}^2(M, N) = \underline{K\text{-mod}}^2(M, N), \quad ev^2: m \otimes f \mapsto (m) f = f(m)$$

$$\underline{K\text{-mod}}^0(M, N) = \underline{K\text{-mod}}^0(M, N), \quad ev^2: f \otimes m \mapsto f(m)$$

Беремо  $L = \underline{K\text{-mod}}^2(M, N)$ ,  $\chi = (L \otimes M \xrightarrow{\text{id}} M \otimes L \xrightarrow{ev^2} N)$

Отримуюмо  $\xi: \underline{K\text{-mod}}^2(M, N) \rightarrow \underline{K\text{-mod}}^2(M, N)$  - канон. ізом.

який дотодужн.  $\text{id}: \underline{K\text{-mod}}^2(M, N) \rightarrow \underline{K\text{-mod}}^2(M, N)$ .

# Градуйовані модулі

Але розглянемо  $gr = Z\text{-grad-}k\text{-mod}$  з тензорним добутком

$$(M \otimes N)^c = \bigoplus_{a+b=c} M^a \otimes N^b \text{ і симетрією } c: M \otimes N \rightarrow N \otimes M;$$

$$m \otimes n \mapsto (-1)^{mn} n \otimes m.$$

$\exists$  градуйовані

$k$ -модулі  $\underline{gr}^a(M, N)$  і  $\underline{gr}^c(M, N)$  разом з паруваннями  $ev^a: M \otimes \underline{gr}^a(M, N) \rightarrow N$

$$ev^c: \underline{gr}^c(M, N) \otimes M \rightarrow N \text{ тжого}$$

$$\forall L \in \dots \quad \forall \varphi: M \otimes L \rightarrow N \exists! \psi: L \rightarrow \underline{gr}^a(M, N);$$

$$\begin{array}{ccc} M \otimes L & & \\ \downarrow \varphi & \searrow \varphi & \\ M \otimes \underline{gr}^a(M, N) & \xrightarrow{ev^a} & N \end{array}$$

$$\forall L \in \dots \quad \forall \chi: L \otimes M \rightarrow N \exists! \xi: L \rightarrow \underline{gr}^c(M, N);$$

$$\begin{array}{ccc} L \otimes M & & \\ \downarrow \xi & \searrow \chi & \\ \underline{gr}^c(M, N) \otimes M & \xrightarrow{ev^c} & N \end{array}$$

# Внутрішні hom'и в градуйованих модулях

Внутрішні hom -  $\underline{gr}^l(M, N)$ ,  $\underline{gr}^r(M, N)$   
 збігаються з  $\underline{gr}(M, N)$  -  $\mathbb{Z}$ -градуйованим  $k$ -модулем з компонентами  
 $\underline{gr}(M, N)^d = \{(f^j : M^j \rightarrow N^{j+d})_{j \in \mathbb{Z}}\}$ , взяттям значення

$$ev^r : m \otimes f \mapsto (m) f = f(m)$$

$$ev^l : f \otimes m \mapsto f(m)$$

Беручи  $\mathcal{L} = \underline{gr}^r \text{Снц} \mathcal{L}$ ,  $\mathcal{X} = (\mathbb{C} \otimes M \xrightarrow{\mathbb{C}} M \otimes \mathbb{C} \xrightarrow{ev^r} N)$

Отримуємо  $\xi : \underline{gr}^r \text{Снц} \mathcal{L} \rightarrow \underline{gr}^l \text{Снц} \mathcal{L}$  - кан. бн. ізом.

Який ототожн. з відображенням

$$\underline{gr}(M, N)^d \rightarrow \underline{gr}(M, N)^d, (f^j)_j \mapsto ((-1)^{jd} f^j)_j.$$

## Диференціально градуйовані модулі = комплекси

розглянемо  $dg = C(k - mod)$  з двома варіантами  $\otimes$  на  $M \otimes N$  у сенсі

$gr$  що відрізняються диференціалом:

$$\text{(лівим)} \quad d(m \otimes n) = (dm) \otimes n + (-1)^m m \otimes (dn) \quad \otimes^l$$

$$\text{(правим)} \quad (m \otimes n)d = m \otimes (nd) + (-1)^n (md) \otimes n \quad \otimes^r$$

Втім, ці моноїдальні категорії ізоморфні завдяки інволютивному функтору

$$\overline{\quad} : dg \rightarrow dg, (M, d) \mapsto \overline{(M, d)} = (M, ((-1)^j d^j)_j). \quad \overline{M \otimes^l N} = \overline{M} \otimes^r \overline{N}.$$

Це замкнена симетрична моноїдальна категорія.

$$\underline{dg}(M, N) = \underline{gr}(M, N) + d$$

З 4х (ізоморфних) варіантів 2:

- ліві диференціали + праві оператори

- праві диференціали + ліві оператори

не дають диференціала на  $\underline{dg}(M, N)$  у вигляді градуйованого комутатора.

Але ліві диференціали + ліві оператори дають диференціал в  $\underline{dg}^l(M, N)$

$$f \mapsto [d, f] = d \circ f - (-1)^f f \circ d$$

праві диференціали + праві оператори дають диференціал в  $\underline{dg}^r(M, N)$

$$f \mapsto [f, d] = f \cdot d - (-1)^f d \cdot f$$

Коланцюгові відображення = циклам в  $\underline{dg}(M, N)^0$

$$f : M \rightarrow N$$

# Гомотопія

$$f \sim 0 \Leftrightarrow \exists s : f = ds.$$

**Definition 1.3.3.** A morphism  $f : X \rightarrow Y$  in  $\mathbf{C}(\mathcal{C})$  is called homotopic to zero if there exist morphisms  $s^n : X^n \rightarrow Y^{n-1}$  in  $\mathcal{C}$  such that for any  $n$ :

$$(1.3.5) \quad f^n = s^{n+1} \circ d_X^n + d_Y^{n-1} \circ s^n .$$

One says  $f$  is **homotopic** to  $g$  if  $f - g$  is homotopic to zero. We denote by  $\text{Ht}(X, Y)$  the subgroup of  $\text{Hom}_{\mathbf{C}(\mathcal{C})}(X, Y)$  consisting of morphisms homotopic to zero. One sees easily that the composition map  $\text{Hom}_{\mathbf{C}(\mathcal{C})}(X, Y) \times \text{Hom}_{\mathbf{C}(\mathcal{C})}(Y, Z) \rightarrow \text{Hom}_{\mathbf{C}(\mathcal{C})}(X, Z)$  sends  $\text{Ht}(X, Y) \times \text{Hom}_{\mathbf{C}(\mathcal{C})}(Y, Z)$  and  $\text{Hom}_{\mathbf{C}(\mathcal{C})}(X, Y) \times \text{Ht}(Y, Z)$  into  $\text{Ht}(X, Z)$ . This permits to define a new category  $\mathbf{K}(\mathcal{C})$  as follows.

**Definition 1.3.4.** The category  $\mathbf{K}(\mathcal{C})$  is defined by

$$(1.3.6) \quad \begin{cases} \text{Ob}(\mathbf{K}(\mathcal{C})) = \text{Ob}(\mathbf{C}(\mathcal{C})) , \\ \text{Hom}_{\mathbf{K}(\mathcal{C})}(X, Y) = \text{Hom}_{\mathbf{C}(\mathcal{C})}(X, Y) / \text{Ht}(X, Y) . \end{cases}$$

One defines similarly the categories  $\mathbf{K}^b(\mathcal{C})$ ,  $\mathbf{K}^+(\mathcal{C})$  and  $\mathbf{K}^-(\mathcal{C})$ . They are full subcategories of  $\mathbf{K}(\mathcal{C})$ .



**Definition 1.3.5.** For  $X \in \text{Ob}(\mathbf{C}(\mathcal{C}))$ , one sets:

$$Z^k(X) = \text{Ker } d_X^k, \quad B^k(X) = \text{Im } d_X^{k-1},$$

$$H^k(X) = \text{Coker}(B^k(X) \rightarrow Z^k(X)).$$

One calls  $H^k(X)$  the  $k$ -th cohomology of the complex  $X$ .

In other words:

$$(1.3.7) \quad H^k(X) = \text{Ker } d_X^k / \text{Im } d_X^{k-1}.$$

Note that  $H^k(\cdot)$  is an additive functor from  $\mathbf{C}(\mathcal{C})$  to  $\mathcal{C}$ , and:

$$(1.3.8) \quad H^k(X) = H^0(X[k]).$$

If  $f : X \rightarrow Y$  is homotopic to zero, then  $H^k(f) : H^k(X) \rightarrow H^k(Y)$  is the zero morphism. Hence  $H^k(\cdot)$  is a well-defined functor from  $\mathbf{K}(\mathcal{C})$  to  $\mathcal{C}$ .

$$\mathbf{K}(\mathbb{k}\text{-mod})(X, Y) = H^0(\underline{\mathbf{dg}}(X, Y)).$$

## Деякі точні послідовності

There are exact sequences:

$$X^{k-1} \longrightarrow Z^k(X) \longrightarrow H^k(X) \longrightarrow 0 ,$$

$$0 \longrightarrow H^k(X) \longrightarrow \text{Coker}(d_X^{k-1}) \longrightarrow X^{k+1} ,$$

$$0 \longrightarrow Z^{k-1}(X) \longrightarrow X^{k-1} \longrightarrow B^k(X) \longrightarrow 0 ,$$

$$0 \longrightarrow B^k(X) \longrightarrow X^k \longrightarrow \text{Coker}(d_X^{k-1}) \longrightarrow 0 ,$$

$$(1.3.9) \quad 0 \rightarrow H^k(X) \rightarrow \text{Coker } d_X^{k-1} \xrightarrow{d_X^k} Z^{k+1}(X) \longrightarrow H^{k+1}(X) \longrightarrow 0 .$$

**Exercise I.9.** Let  $\mathcal{C}$  be an abelian category. Consider the commutative diagram with exact rows in  $\mathcal{C}$ :

$$\begin{array}{ccccccc}
 & & X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \longrightarrow & 0 \\
 & & \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \\
 0 & \longrightarrow & X' & \xrightarrow{f'} & Y' & \xrightarrow{g'} & Z' & & 
 \end{array}$$

Prove that there is a natural exact sequence:

$$\text{Ker } \alpha \longrightarrow \text{Ker } \beta \longrightarrow \text{Ker } \gamma \xrightarrow{\varphi} \text{Coker } \alpha \longrightarrow \text{Coker } \beta \longrightarrow \text{Coker } \gamma ,$$

so that the following diagram commutes:

$$\begin{array}{ccccc}
 & & Y & \xrightarrow{g} & Z \\
 & & \uparrow & & \uparrow \\
 Y & \longleftarrow & \text{Ker } \gamma \circ g & \longrightarrow & \text{Ker } \gamma \\
 \downarrow & & \downarrow & & \downarrow \varphi \\
 Y' & \xleftarrow{f'} & X' & \longrightarrow & \text{Coker } \alpha .
 \end{array}$$

## Довга точна послідовність

**Proposition 1.3.6.** Let  $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$  be an exact sequence in  $\mathbf{C}(\mathcal{C})$ . Then there exists a canonical long exact sequence in  $\mathcal{C}$ :

$$\cdots \longrightarrow H^n(X) \longrightarrow H^n(Y) \longrightarrow H^n(Z) \xrightarrow{\delta} H^{n+1}(X) \longrightarrow \cdots,$$

more precisely, if

$$\begin{array}{ccccccccc} 0 & \longrightarrow & X & \longrightarrow & Y & \longrightarrow & Z & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & X' & \longrightarrow & Y' & \longrightarrow & Z' & \longrightarrow & 0 \end{array}$$

is a commutative diagram of exact sequences in  $\mathbf{C}(\mathcal{C})$ , then all the diagrams:

$$\begin{array}{ccc} H^n(Z) & \longrightarrow & H^{n+1}(X) \\ \downarrow & & \downarrow \\ H^n(Z') & \longrightarrow & H^{n+1}(X') \end{array}$$

commute.

## Доведення довгої точної послідовності

*Proof.* Consider the commutative diagram with exact rows:

$$\begin{array}{ccccccc} \text{Coker}(d_X^{n-1}) & \longrightarrow & \text{Coker}(d_Y^{n-1}) & \longrightarrow & \text{Coker}(d_Z^{n-1}) & \longrightarrow & 0 \\ \downarrow d_X^n & & \downarrow d_Y^n & & \downarrow d_Z^n & & \\ 0 & \longrightarrow & Z^{n+1}(X) & \longrightarrow & Z^{n+1}(Y) & \longrightarrow & Z^{n+1}(Z) . \end{array}$$

The result then follows from (1.3.9) and Exercise I.9. The functoriality of the construction is left to the reader.  $\square$

# Гомотопічні ізоморфізми. Ланцюгові комплекси. Гомології

**Remark 1.3.8.** Let  $X$  and  $Y$  be two objects of  $\mathbf{C}(\mathcal{C})$ . One sometimes says that  $X$  and  $Y$  are homotopically equivalent if they are isomorphic in  $\mathbf{K}(\mathcal{C})$ , that is, if there exists  $f \in \text{Hom}_{\mathbf{C}(\mathcal{C})}(X, Y)$  which is an isomorphism in  $\mathbf{K}(\mathcal{C})$ . Such an  $f$  is called a homotopy equivalence.

**Notations 1.3.9.** (i) Consider a sequence  $\{X_n, d_n^X\}_{n \in \mathbb{Z}}$  where  $X_n \in \text{Ob}(\mathcal{C})$ ,  $d_n^X \in \text{Hom}_{\mathcal{C}}(X_n, X_{n-1})$  and  $d_n^X \circ d_{n+1}^X = 0$ . Then we shall still say that this sequence is a complex in  $\mathcal{C}$ . In fact setting  $X^n = X_{-n}$ ,  $d_X^n = d_{-n}^X$ , the sequence  $\{X^n, d_X^n\}$  is a complex in our previous sense.

(ii) We sometimes denote by  $X^\cdot$  (resp.  $X_*$ ) a complex  $\{X^n, d_X^n\}$  (resp.  $\{X_n, d_n^X\}$ ). The object  $\text{Ker } d_{n-1}^X / \text{Im } d_n^X$  is called the  $n$ -th homology group of  $X$ , and denoted by  $H_n(X)$ .

## Конус відображення

Let  $\mathcal{C}$  be an additive category, and let  $f : X \rightarrow Y$  be a morphism in  $\mathbf{C}(\mathcal{C})$ .

**Definition 1.4.1.** *The mapping cone of  $f$ , denoted by  $M(f)$ , is the object of  $\mathbf{C}(\mathcal{C})$  defined as follows:*

$$(1.4.1) \quad \begin{cases} M(f)^n = X^{n+1} \oplus Y^n, \\ d_{M(f)}^n = \begin{pmatrix} d_{X[1]}^n & 0 \\ f^{n+1} & d_Y^n \end{pmatrix}. \end{cases}$$

Recall that  $d_{X[1]}^n = -d_X^{n+1}$ .

We define the morphisms  $\alpha(f) : Y \rightarrow M(f)$  and  $\beta(f) : M(f) \rightarrow X[1]$  by:

$$(1.4.2) \quad \alpha(f)^n = \begin{pmatrix} 0 \\ \text{id}_{Y^n} \end{pmatrix},$$

$$(1.4.3) \quad \beta(f)^n = (\text{id}_{X^{n+1}}, 0).$$

In other notation  $d_{M(f)} = \begin{pmatrix} d[1] & 0 \\ f \circ \sigma^{-1} & d \end{pmatrix}$ .

## Перше обертання трикутника

**Lemma 1.4.2.** For any  $f : X \rightarrow Y$  in  $\mathbf{C}(\mathcal{C})$ , there exists  $\phi : X[1] \rightarrow M(\alpha(f))$  such that:

(1.4.4)  $\phi$  is an isomorphism in  $\mathbf{K}(\mathcal{C})$ ,

(1.4.5) The diagram below commutes in  $\mathbf{K}(\mathcal{C})$ :

$$\begin{array}{ccccccc} Y & \xrightarrow{\alpha(f)} & M(f) & \xrightarrow{\beta(f)} & X[1] & \xrightarrow{-f[1]} & Y[1] \\ \downarrow \text{id}_Y & & \downarrow \text{id}_{M(f)} & & \downarrow \phi & & \downarrow \text{id}_{Y[1]} \\ Y & \xrightarrow{\alpha(f)} & M(f) & \xrightarrow{\alpha(\alpha(f))} & M(\alpha(f)) & \xrightarrow{\beta(\alpha(f))} & Y[1] . \end{array}$$

Note that such a result would not hold in  $\mathbf{C}(\mathcal{C})$ . Note further that  $\phi$  is not unique



*Proof.* We have:

$$M(\alpha(f))^n = Y^{n+1} \oplus M(f)^n = Y^{n+1} \oplus X^{n+1} \oplus Y^n .$$

We define  $\phi^n : X[1]^n \rightarrow M(\alpha(f))^n$  and  $\psi^n : M(\alpha(f))^n \rightarrow X[1]^n$  by:

$$\phi^n = \begin{pmatrix} -f^{n+1} \\ \text{id}_{X^{n+1}} \\ 0 \end{pmatrix}, \quad \psi^n = (0, \text{id}_{X^{n+1}}, 0) .$$

Then the lemma follows from the following observations.

- (a)  $\phi = (\phi^n)_n$  and  $\psi = (\psi^n)_n$  are morphisms of complexes,
- (b)  $\psi \circ \phi = \text{id}_{X[1]}$ ,
- (c)  $\phi \circ \psi$  is homotopic to  $\text{id}_{M(\alpha(f))}$ ,
- (d)  $\psi \circ \alpha(\alpha(f)) = \beta(f)$ ,
- (e)  $\beta(\alpha(f)) \circ \phi = -f[1]$ .

All these properties, except (c), can be checked directly. To get (c) we define  $s^n : M(\alpha(f))^n \rightarrow M(\alpha(f))^{n-1}$  by:

$$s^n = \begin{pmatrix} 0 & 0 & \text{id}_{Y^n} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} .$$

Then one verifies that:

$$\text{id}_{M(\alpha(f))^n} - \phi^n \circ \psi^n = s^{n+1} \circ d_{M(\alpha(f))}^n + d_{M(\alpha(f))}^{n-1} \circ s^n . \quad \square$$

$$M(\alpha(f)) = Y[1] \oplus X[1] \oplus Y, \quad d_{M(\alpha(f))} = \begin{pmatrix} d[1] & 0 & 0 \\ 0 & d[1] & 0 \\ \sigma^{-1} & f \circ \sigma^{-1} & d \end{pmatrix}.$$

$$\phi = \begin{pmatrix} -f[1] \\ 1 \\ 0 \end{pmatrix}, \quad \psi = (0 \ 1 \ 0), \quad s = \begin{pmatrix} 0 & 0 & \sigma \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

(c)

$$\begin{aligned} 1_{M(\alpha(f))} - \phi \circ \psi &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 0 & -f[1] & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & f[1] & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 0 & 0 & \sigma \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} d[1] & 0 & 0 \\ 0 & d[1] & 0 \\ \sigma^{-1} & f \circ \sigma^{-1} & d \end{pmatrix} \\ &+ \begin{pmatrix} d[1] & 0 & 0 \\ 0 & d[1] & 0 \\ \sigma^{-1} & f \circ \sigma^{-1} & d \end{pmatrix} \begin{pmatrix} 0 & 0 & \sigma \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = s \circ d + d \circ s \end{aligned}$$

завдяки  $\sigma \circ d + d[1] \circ \sigma = 0$ .

## Виділені трикутники

One defines a **triangle** in  $\mathbf{K}(\mathcal{C})$  as being a sequence of morphisms  $X \rightarrow Y \rightarrow Z \rightarrow X[1]$  and a morphism of triangles as being a commutative diagram in  $\mathbf{K}(\mathcal{C})$ :

$$\begin{array}{ccccccc} X & \longrightarrow & Y & \longrightarrow & Z & \longrightarrow & X[1] \\ \downarrow \phi & & \downarrow & & \downarrow & & \downarrow \phi[1] \\ X' & \longrightarrow & Y' & \longrightarrow & Z' & \longrightarrow & X'[1] . \end{array}$$

**Definition 1.4.3.** A triangle  $X \rightarrow Y \rightarrow Z \rightarrow X[1]$  in  $\mathbf{K}(\mathcal{C})$  is called a distinguished triangle, if it is isomorphic to a triangle  $X' \xrightarrow{f} Y' \xrightarrow{\alpha(f)} M(f) \xrightarrow{\beta(f)} X'[1]$ , for some  $f$  in  $\mathbf{C}(\mathcal{C})$ .

# Властивості $\mathbf{K}(\mathcal{C})$ як триангульованої категорії

**Proposition 1.4.4.** *The collection of distinguished triangles in  $\mathbf{K}(\mathcal{C})$  satisfies the following properties, (TR 0)–(TR 5).*

(TR 0) *A triangle isomorphic to a distinguished triangle is distinguished.*

(TR 1) *For any  $X \in \text{Ob}(\mathbf{K}(\mathcal{C}))$ ,  $X \xrightarrow{\text{id}_X} X \longrightarrow 0 \longrightarrow X[1]$  is a distinguished triangle.*

(TR 2) *Any  $f : X \rightarrow Y$  in  $\mathbf{K}(\mathcal{C})$  can be embedded in a distinguished triangle  $X \xrightarrow{f} Y \rightarrow Z \rightarrow X[1]$ .*

(TR 3)  *$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} X[1]$  is a distinguished triangle if and only if  $Y \xrightarrow{g} Z \xrightarrow{h} X[1] \xrightarrow{-f[1]} Y[1]$  is a distinguished triangle.*

(TR 4) *Given two distinguished triangles  $X \xrightarrow{f} Y \rightarrow Z \rightarrow X[1]$  and  $X' \xrightarrow{f'} Y' \rightarrow Z' \rightarrow X'[1]$ , a commutative diagram*

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow u & & \downarrow v \\ X' & \xrightarrow{f'} & Y' \end{array}$$

*can be embedded in a morphism of triangles (not necessarily unique).*

(TR 5) (octahedral axiom). *Suppose given distinguished triangles:*

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \longrightarrow & Z' & \longrightarrow & X[1] , \\ Y & \xrightarrow{g} & Z & \longrightarrow & X' & \longrightarrow & Y[1] , \\ X & \xrightarrow{g \circ f} & Z & \longrightarrow & Y' & \longrightarrow & X[1] , \end{array}$$

*then there exists a distinguished triangle*

$$Z' \rightarrow Y' \rightarrow X' \rightarrow Z'[1]$$

*such that the following diagram is commutative:*

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \longrightarrow & Z' & \longrightarrow & X[1] \\ \text{id}_X \downarrow & & \downarrow g & & \downarrow & & \downarrow \text{id}_{X[1]} \\ X & \xrightarrow{g \circ f} & Z & \longrightarrow & Y' & \longrightarrow & X[1] \\ f \downarrow & & \downarrow \text{id}_Z & & \downarrow & & \downarrow f[1] \\ Y & \xrightarrow{g} & Z & \longrightarrow & X' & \longrightarrow & Y[1] \\ \downarrow & & \downarrow & & \downarrow \text{id}_{X'} & & \downarrow \\ Z' & \longrightarrow & Y' & \longrightarrow & X' & \longrightarrow & Z'[1] . \end{array}$$

## Доведення властивостей $\mathbf{K}(\mathcal{C})$ як трианг. категорії

*Proof.* The properties (TR 0) and (TR 2) are obvious, and (TR 3) follows from Lemma 1.4.2.

Since the mapping cone of  $f : 0 \rightarrow X$  is  $X$ , the triangle  $0 \rightarrow X \xrightarrow{\text{id}_X} X \rightarrow 0[1]$  is distinguished. Applying (TR 3) we get (TR 1). Let us prove (TR 4). We may assume that  $X \xrightarrow{f} Y \rightarrow Z \rightarrow X[1]$  and  $X' \xrightarrow{f'} Y' \rightarrow Z' \rightarrow X'[1]$  are  $X \xrightarrow{f} Y \xrightarrow{\alpha(f)} M(f) \xrightarrow{\beta(f)} X[1]$  and  $X' \xrightarrow{f'} Y' \xrightarrow{\alpha(f')} M(f') \xrightarrow{\beta(f')} X'[1]$ , respectively. We shall construct a morphism  $w : M(f) \rightarrow M(f')$  such that:

$$(1.4.4) \quad \begin{cases} w \circ \alpha(f) = \alpha(f') \circ v , \\ u[1] \circ \beta(f) = \beta(f') \circ w . \end{cases}$$

By the definition of  $\mathbf{K}(\mathcal{C})$ , there exists  $s^n : X^n \rightarrow Y'^{n-1}$  such that  $v^n \circ f^n - f'^n \circ u^n = s^{n+1} \circ d_X^n + d_{Y'}^{n-1} \circ s^n$ . We define  $w^n : M(f)^n = X^{n+1} \oplus Y^n \rightarrow M(f')^n = X'^{n+1} \oplus Y'^n$  by:

$$w^n = \begin{pmatrix} u^{n+1} & 0 \\ s^{n+1} & v^n \end{pmatrix} .$$

Then a direct calculation shows that  $w$  is a morphism of complexes and satisfies (1.4.4).

Let us prove (TR 5). We may assume  $Z' = M(f)$ ,  $X' = M(g)$  and  $Y' = M(g \circ f)$ . Let us define  $u: Z' \rightarrow Y'$  and  $v: Y' \rightarrow X'$  by:

$$u^n: X^{n+1} \oplus Y^n \rightarrow X^{n+1} \oplus Z^n, \quad u = \begin{pmatrix} \text{id}_{X^{n+1}} & 0 \\ 0 & g^n \end{pmatrix},$$

$$v^n: X^{n+1} \oplus Z^n \rightarrow Y^{n+1} \oplus Z^n, \quad v^n = \begin{pmatrix} f^{n+1} & 0 \\ 0 & \text{id}_{Z^n} \end{pmatrix}.$$

We define  $w: X' \rightarrow Z'[1]$  as the composite  $X' \rightarrow Y[1] \rightarrow Z'[1]$ . Then the diagram in (TR 5) is commutative, and it is enough to show that  $Z' \xrightarrow{u} Y' \xrightarrow{v} X' \xrightarrow{w} Z'[1]$  is a distinguished triangle. For that purpose we shall construct an isomorphism  $\phi: M(u) \rightarrow X'$  and its inverse  $\psi: X' \rightarrow M(u)$  such that  $\phi \circ \alpha(u) = v$  and  $\beta(u) \circ \psi = w$ . We have:

$$M(u)^n = M(f)^{n+1} \oplus M(g \circ f)^n = X^{n+2} \oplus Y^{n+1} \oplus X^{n+1} \oplus Z^n$$

and  $X'^n = M(g)^n = Y^{n+1} \oplus Z^n$ . We define  $\phi$  and  $\psi$  by:

$$\phi^n = \begin{pmatrix} 0 & \text{id}_{Y^{n+1}} & f^{n+1} & 0 \\ 0 & 0 & 0 & \text{id}_{Z^n} \end{pmatrix}, \quad \psi^n = \begin{pmatrix} 0 & 0 \\ \text{id}_{Y^{n+1}} & 0 \\ 0 & 0 \\ 0 & \text{id}_{X^{n+1}} \end{pmatrix}.$$

Then one checks easily that  $\phi$  and  $\psi$  are morphisms of complexes and  $\phi \circ \alpha(u) = v$ ,  $\beta(u) \circ \psi = w$ .

# Октаэдр

We have  $\phi \circ \psi = \text{id}_{X'}$ . If we define:


$$s^n : M(u)^n \rightarrow M(u)^{n-1} , \quad s^n = \begin{pmatrix} 0 & 0 & \text{id}_{X^{n+1}} & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

then:

$$(\text{id}_{M(u)} - \psi \circ \phi)^n = s^{n+1} \circ d_{M(u)}^n + d_{M(u)}^{n-1} \circ s^n .$$

Hence  $\psi \circ \phi$  equals  $\text{id}_{M(u)}$  in  $\mathbf{K}(\mathcal{C})$ .  $\square$

**Remark 1.4.5.** Property (TR 5) may be visualized by the following octahedral diagram:







Masaki Kashiwara and Pierre Schapira, Sheaves on manifolds, Grundlehren der mathematischen Wissenschaften, vol. 292, Springer-Verlag, Berlin, New York, 1990. §1.3–§1.4