

7. Породження триангульованих категорій. Навколо похідних категорій

Володимир Любашенко

18 березня 2021

Let us recall that when \mathcal{A} is a small additive category, then $K(\mathcal{A})$ denotes the homotopy category of complexes. Namely, its objects are cochain complexes of objects in \mathcal{A} , while its morphisms are homotopy equivalence classes of morphisms of complexes. For $A^* \in \text{Ob}(K(\mathcal{A}))$, we denote by A^i its i -th component. We can then define the full subcategories $K^b(\mathcal{A})$, $K^+(\mathcal{A})$, $K^-(\mathcal{A})$ of the category $K(\mathcal{A})$ whose objects are

$$\text{Ob}(K^b(\mathcal{A})) = \{A^* \in K(\mathcal{A}) \mid A^i = 0 \text{ for all } |i| \gg 0\}$$

$$\text{Ob}(K^+(\mathcal{A})) = \{A^* \in K(\mathcal{A}) \mid A^i = 0 \text{ for all } i \ll 0\}$$

$$\text{Ob}(K^-(\mathcal{A})) = \{A^* \in K(\mathcal{A}) \mid A^i = 0 \text{ for all } i \gg 0\}$$

Let us recall that when \mathcal{A} is a small additive category, then $K(\mathcal{A})$ denotes the homotopy category of complexes. Namely, its objects are cochain complexes of objects in \mathcal{A} , while its morphisms are homotopy equivalence classes of morphisms of complexes. For $A^* \in \text{Ob}(K(\mathcal{A}))$, we denote by A^i its i -th component. We can then define the full subcategories $K^b(\mathcal{A})$, $K^+(\mathcal{A})$, $K^-(\mathcal{A})$ of the category $K(\mathcal{A})$ whose objects are

$$\text{Ob}(K^b(\mathcal{A})) = \{A^* \in K(\mathcal{A}) \mid A^i = 0 \text{ for all } |i| \gg 0\}$$

$$\text{Ob}(K^+(\mathcal{A})) = \{A^* \in K(\mathcal{A}) \mid A^i = 0 \text{ for all } i \ll 0\}$$

$$\text{Ob}(K^-(\mathcal{A})) = \{A^* \in K(\mathcal{A}) \mid A^i = 0 \text{ for all } i \gg 0\}$$

For $? = b, +, -, \emptyset$, we single out the full subcategory $V^?(\mathcal{A}) \subseteq K^?(\mathcal{A})$ consisting of objects with zero differentials. It will be crucial in the rest. Here we just point out that, for an object $A^* \in V^?(\mathcal{A})$, we will use the shorthand

$$\bigoplus_{i \in \mathbb{Z}} A^i[-i]$$

to remind that the object $A^i \in \mathcal{A}$ is placed in degree i .

When \mathcal{A} is an abelian category, the full triangulated subcategory $K_{\text{acy}}^?(\mathcal{A}) \subseteq K^?(\mathcal{A})$ consists of acyclic complexes, i.e. objects in $K(\mathcal{A})$ with trivial cohomology.

When \mathcal{A} is an abelian category, the full triangulated subcategory $K_{\text{acy}}^?(\mathcal{A}) \subseteq K^?(\mathcal{A})$ consists of acyclic complexes, i.e. objects in $K(\mathcal{A})$ with trivial cohomology.

The triangulated category $D^?(\mathcal{A})$ is then the Verdier quotient of $K^?(\mathcal{A})$ by $K_{\text{acy}}^?(\mathcal{A})$, and it comes with a quotient functor

$$Q : K^?(\mathcal{A}) \longrightarrow D^?(\mathcal{A}).$$

When \mathcal{A} is an abelian category, the full triangulated subcategory $K_{\text{acy}}^?(\mathcal{A}) \subseteq K^?(\mathcal{A})$ consists of acyclic complexes, i.e. objects in $K(\mathcal{A})$ with trivial cohomology.

The triangulated category $D^?(\mathcal{A})$ is then the Verdier quotient of $K^?(\mathcal{A})$ by $K_{\text{acy}}^?(\mathcal{A})$, and it comes with a quotient functor

$$Q : K^?(\mathcal{A}) \longrightarrow D^?(\mathcal{A}).$$

We can then consider the full subcategory $B^?(\mathcal{A}) \subseteq D^?(\mathcal{A})$ as

$$B^?(\mathcal{A}) := Q(V^?(\mathcal{A})).$$

Породження триангульованих категорій

Definition

Let \mathcal{T} be a triangulated category and let $\mathcal{S} \subset \mathbf{Ob}(\mathcal{T})$. We define

1. $\langle \mathcal{S} \rangle_1$ is the collection of all direct summands of finite coproducts of shifts of objects in \mathcal{S} ;

Породження триангульованих категорій

Definition

Let \mathcal{T} be a triangulated category and let $\mathcal{S} \subset \text{Ob}(\mathcal{T})$. We define

1. $\langle \mathcal{S} \rangle_1$ is the collection of all direct summands of finite coproducts of shifts of objects in \mathcal{S} ;
2. $\langle \mathcal{S} \rangle_{n+1}$ consists of all direct summands of objects $T \in \mathcal{T}$, for which there exists a distinguished triangle $T_1 \rightarrow T \rightarrow T_2$ with $T_1 \in \langle \mathcal{S} \rangle_n$ and $T_2 \in \langle \mathcal{S} \rangle_1$.

Породження триангульованих категорій

Definition

Let \mathcal{T} be a triangulated category and let $\mathcal{S} \subset \text{Ob}(\mathcal{T})$. We define

1. $\langle \mathcal{S} \rangle_1$ is the collection of all direct summands of finite coproducts of shifts of objects in \mathcal{S} ;
2. $\langle \mathcal{S} \rangle_{n+1}$ consists of all direct summands of objects $T \in \mathcal{T}$, for which there exists a distinguished triangle $T_1 \rightarrow T \rightarrow T_2$ with $T_1 \in \langle \mathcal{S} \rangle_n$ and $T_2 \in \langle \mathcal{S} \rangle_1$.

We set $\langle \mathcal{S} \rangle_\infty$ for the full subcategory consisting of all objects T in \mathcal{T} contained in $\langle \mathcal{S} \rangle_n$, for some n .

Proposition

Recall $V^?(A) \subset K^?(A)$. For $? = b, +, -, \emptyset$, we have that

$$\langle V^?(A) \rangle_3 = K^?(A).$$

Proposition

Recall $V^?(A) \subset K^?(A)$. For $? = b, +, -, \emptyset$, we have that

$$\langle V^?(A) \rangle_3 = K^?(A).$$

Доведення. Let $A^* \in \text{Ob}(K^?(A))$, which we write as a complex

$$\dots \longrightarrow A^{-2} \longrightarrow A^{-1} \longrightarrow A^0 \longrightarrow A^1 \longrightarrow A^2 \longrightarrow \dots$$

Let K^i be the kernel of the differential $A^i \rightarrow A^{i+1}$. Then the map $A^{i-1} \rightarrow A^i$ factors uniquely as $A^{i-1} \xrightarrow{\alpha^i} K^i \hookrightarrow A^i$.

Proposition

Recall $V^?(A) \subset K^?(A)$. For $? = b, +, -, \emptyset$, we have that $\langle V^?(A) \rangle_3 = K^?(A)$.

Доведення. Let $A^* \in \text{Ob}(K^?(A))$, which we write as a complex

$$\dots \longrightarrow A^{-2} \longrightarrow A^{-1} \longrightarrow A^0 \longrightarrow A^1 \longrightarrow A^2 \longrightarrow \dots$$

Let K^i be the kernel of the differential $A^i \rightarrow A^{i+1}$. Then the map $A^{i-1} \rightarrow A^i$ factors uniquely as $A^{i-1} \xrightarrow{\alpha^i} K^i \hookrightarrow A^i$.

This yields the morphism

$$\bigoplus_{i \in \mathbb{Z}} A^{i-1}[-i] \xrightarrow{\bigoplus_{i \in \mathbb{Z}} \alpha^i} \bigoplus_{i \in \mathbb{Z}} K^i[-i]$$

in $V^?(A)$. Denote by C^* its mapping cone. It is clear that $C^* \in \langle V^?(A) \rangle_2$ and it is the direct sum over $i \in \mathbb{Z}$ of the complexes

$$\dots \longrightarrow 0 \longrightarrow A^{i-1} \xrightarrow{\alpha^i} K^i \longrightarrow 0 \longrightarrow \dots$$

Now consider the cochain map

$$\bigoplus_{i \in \mathbb{Z}} K^i[-i] \xrightarrow{\varphi + \psi} C^*$$

whose components, out of $K^i[-i]$, are (respectively) φ^i as below

$$\begin{array}{ccccccccccc} \dots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & K^i & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \dots \\ & & \downarrow & & \downarrow & & \parallel & & \downarrow & & \downarrow & & \\ \dots & \longrightarrow & 0 & \longrightarrow & A^{i-1} & \longrightarrow & K^i & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \dots \end{array}$$

Now consider the cochain map

$$\bigoplus_{i \in \mathbb{Z}} K^i[-i] \xrightarrow{\varphi + \psi} C^*$$

whose components, out of $K^i[-i]$, are (respectively) φ^i as below

$$\begin{array}{ccccccccccc} \dots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & K^i & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \dots \\ & & \downarrow & & \downarrow & & \parallel & & \downarrow & & \downarrow & & \\ \dots & \longrightarrow & 0 & \longrightarrow & A^{i-1} & \longrightarrow & K^i & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \dots \end{array}$$

and ψ^i as below

$$\begin{array}{ccccccccccc} \dots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & K^i & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \dots \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\ \dots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & A^i & \longrightarrow & K^{i+1} & \longrightarrow & 0 & \longrightarrow & \dots \end{array}$$

Породження похідної категорії в 3 кроки

It can be easily checked that the mapping cone of the morphism $\varphi + \psi$ is isomorphic to the direct sum of the complex A^* and of complexes of the form

$$\dots \longrightarrow 0 \longrightarrow K^i \longrightarrow K^i \longrightarrow 0 \longrightarrow \dots$$

Породження похідної категорії в 3 кроки

It can be easily checked that the mapping cone of the morphism $\varphi + \psi$ is isomorphic to the direct sum of the complex A^* and of complexes of the form

$$\dots \longrightarrow 0 \longrightarrow K^i \longrightarrow K^i \longrightarrow 0 \longrightarrow \dots$$

In other words, $\text{Cone}(\varphi + \psi) \cong A^*$ in $K^?(\mathcal{A})$. Therefore, as C^* belongs to $\langle V^?(\mathcal{A}) \rangle_2$ and $\varphi + \psi$ is a morphism from an object of $V^?(\mathcal{A})$ to C^* , we have that $A^* \in \langle V^?(\mathcal{A}) \rangle_3$. \square

Породження похідної категорії в 3 кроки

It can be easily checked that the mapping cone of the morphism $\varphi + \psi$ is isomorphic to the direct sum of the complex A^* and of complexes of the form

$$\dots \longrightarrow 0 \longrightarrow K^i \longrightarrow K^i \longrightarrow 0 \longrightarrow \dots$$

In other words, $\text{Cone}(\varphi + \psi) \cong A^*$ in $K^?(A)$. Therefore, as C^* belongs to $\langle V^?(A) \rangle_2$ and $\varphi + \psi$ is a morphism from an object of $V^?(A)$ to C^* , we have that $A^* \in \langle V^?(A) \rangle_3$. \square

Corollary

Recall $B^?(A) \subset D^?(A)$. For $? = b, +, -, \emptyset$, we have that $\langle B^?(A) \rangle_3 = D^?(A)$.

Cone $(\varphi + \psi)^i = (K^{i+1} \oplus A^i \oplus K^i, d)$ has the differential

$$d = \begin{pmatrix} 0 & \iota^{i+1} & 1 \\ 0 & 0 & \alpha^{i+1} \\ 0 & 0 & 0 \end{pmatrix}. \text{ We have}$$

$$(A^{i-1} \xrightarrow{\alpha^i} K^i \xrightarrow{\iota^i} A^i) = d, \quad (K^i \xrightarrow{\iota^i} A^i \xrightarrow{\alpha^{i+1}} K^{i+1}) = 0.$$

Cone $(\varphi + \psi)^i = (K^{i+1} \oplus A^i \oplus K^i, d)$ has the differential

$$d = \begin{pmatrix} 0 & \iota^{i+1} & 1 \\ 0 & 0 & \alpha^{i+1} \\ 0 & 0 & 0 \end{pmatrix}. \text{ We have}$$

$$(A^{i-1} \xrightarrow{\alpha^i} K^i \xrightarrow{\iota^i} A^i) = d, \quad (K^i \xrightarrow{\iota^i} A^i \xrightarrow{\alpha^{i+1}} K^{i+1}) = 0.$$

There is a chain map

$$\begin{array}{ccc} \rightarrow K^{i+1} \oplus A^i \oplus K^i & \xrightarrow{\begin{pmatrix} 0 & \iota^{i+1} & 1 \\ 0 & 0 & \alpha^{i+1} \\ 0 & 0 & 0 \end{pmatrix}} & K^{i+2} \oplus A^{i+1} \oplus K^{i+1} \rightarrow \\ \begin{pmatrix} 1 & 0 & 0 \\ \alpha^{i+1} & 1 & 0 \\ 0 & -\iota^i & 1 \end{pmatrix} \downarrow & & \downarrow \begin{pmatrix} 1 & 0 & 0 \\ \alpha^{i+2} & 1 & 0 \\ 0 & -\iota^{i+1} & 1 \end{pmatrix} \\ \rightarrow K^{i+1} \oplus A^i \oplus K^i & \xrightarrow{\begin{pmatrix} 0 & 0 & 1 \\ 0 & -d & 0 \\ 0 & 0 & 0 \end{pmatrix}} & K^{i+2} \oplus A^{i+1} \oplus K^{i+1} \rightarrow \end{array}$$

Cone $(\varphi + \psi)^i = (K^{i+1} \oplus A^i \oplus K^i, d)$ has the differential

$$d = \begin{pmatrix} 0 & \iota^{i+1} & 1 \\ 0 & 0 & \alpha^{i+1} \\ 0 & 0 & 0 \end{pmatrix}. \text{ We have}$$

$$(A^{i-1} \xrightarrow{\alpha^i} K^i \xrightarrow{\iota^i} A^i) = d, \quad (K^i \xrightarrow{\iota^i} A^i \xrightarrow{\alpha^{i+1}} K^{i+1}) = 0.$$

There is a chain map

$$\begin{array}{ccc} \rightarrow K^{i+1} \oplus A^i \oplus K^i & \xrightarrow{\begin{pmatrix} 0 & \iota^{i+1} & 1 \\ 0 & 0 & \alpha^{i+1} \\ 0 & 0 & 0 \end{pmatrix}} & K^{i+2} \oplus A^{i+1} \oplus K^{i+1} \rightarrow \\ \begin{pmatrix} 1 & 0 & 0 \\ \alpha^{i+1} & 1 & 0 \\ 0 & -\iota^i & 1 \end{pmatrix} \downarrow & & \downarrow \begin{pmatrix} 1 & 0 & 0 \\ \alpha^{i+2} & 1 & 0 \\ 0 & -\iota^{i+1} & 1 \end{pmatrix} \\ \rightarrow K^{i+1} \oplus A^i \oplus K^i & \xrightarrow{\begin{pmatrix} 0 & 0 & 1 \\ 0 & -d & 0 \\ 0 & 0 & 0 \end{pmatrix}} & K^{i+2} \oplus A^{i+1} \oplus K^{i+1} \rightarrow \end{array}$$

It is invertible $\Rightarrow \text{Cone}(\varphi + \psi) \cong ((A^i), -d) \in K^?(\mathcal{A})$.

Модельна структура на \mathbf{dgCat}

If we consider for \mathcal{C} the category \mathbf{DCAT} , for W the subcategory of quasi-equivalences, the category \mathbf{DCAT} admits a Quillen model structure whose weak equivalences are the quasi-equivalences.

Модельна структура на \mathbf{dgCat}

If we consider for \mathcal{C} the category \mathbf{DCAT} , for W the subcategory of quasi-equivalences, the category \mathbf{DCAT} admits a Quillen model structure whose weak equivalences are the quasi-equivalences.

\mathbf{dgCat} has a model structure whose weak equivalences are quasi-equivalences and such that every object is fibrant. We denote by \mathbf{Hqe} the corresponding homotopy category, namely the localization of \mathbf{dgCat} with respect to quasi-equivalences. Since H^0 sends quasi-equivalences to equivalences, for every morphism $f: \mathcal{C}_1 \rightarrow \mathcal{C}_2$ in \mathbf{Hqe} there is a \mathbb{k} -linear functor $H^0(f): H^0(\mathcal{C}_1) \rightarrow H^0(\mathcal{C}_2)$, which is well-defined up to equivalences.

Модельна структура на \mathbf{dgCat}

If we consider for \mathcal{C} the category \mathbf{DCAT} , for W the subcategory of quasi-equivalences, the category \mathbf{DCAT} admits a Quillen model structure whose weak equivalences are the quasi-equivalences.

\mathbf{dgCat} has a model structure whose weak equivalences are quasi-equivalences and such that every object is fibrant. We denote by \mathbf{Hqe} the corresponding homotopy category, namely the localization of \mathbf{dgCat} with respect to quasi-equivalences. Since H^0 sends quasi-equivalences to equivalences, for every morphism $f: \mathcal{C}_1 \rightarrow \mathcal{C}_2$ in \mathbf{Hqe} there is a \mathbb{k} -linear functor $H^0(f): H^0(\mathcal{C}_1) \rightarrow H^0(\mathcal{C}_2)$, which is well-defined up to equivalences.

Dg functors between two dg categories \mathcal{C}_1 and \mathcal{C}_2 form in a natural way the objects of a dg category $\underline{\mathbf{Hom}}(\mathcal{C}_1, \mathcal{C}_2)$. For every dg category \mathcal{C} we set $\mathbf{dgMod}(\mathcal{C}) := \underline{\mathbf{Hom}}(\mathcal{C}^{\text{op}}, \mathbf{C}_{\text{dg}}(\text{Mod}(\mathbb{k})))$ and call its objects (right) dg \mathcal{C} -modules.

Зсуви в dg-категорії

Given an object $A \in \mathcal{A}$, the object $A[r]$ is characterized (up to a DG isomorphism) by the existence of closed morphisms $f : A \rightarrow A[r]$, $g : A[r] \rightarrow A$ of degrees $-r$ and r , respectively, such that $fg = gf = 1$. Thus, in particular, every DG functor commutes with shifts.

Definition 4.6. The objects of $\mathcal{A}^{\text{pre-tr}}$ are “one-sided twisted complexes,” that is, formal expressions $(\oplus_{i=1}^n C_i[r_i], q)$, where $C_i \in \text{Ob } \mathcal{A}$, $r_i \in \mathbb{Z}$, $n \geq 0$, $q = (q_{ij})$, $q_{ij} \in \text{Hom}(C_j[r_j], C_i[r_i])$ is homogeneous of degree 1, $q_{ij} = 0$ for $i \geq j$, and $dq + q^2 = 0$. If $C, C' \in \text{Ob } \mathcal{A}^{\text{pre-tr}}$, $C = (\oplus C_j[r_j], q)$, $C' = (\oplus C'_j[r'_j], q')$, then the \mathbb{Z} -graded k -module $\text{Hom}(C, C')$ is the space of matrices $f = (f_{ij})$, $f_{ij} \in \text{Hom}(C_j[r_j], C'_i[r'_i])$, and the composition map $\text{Hom}(C, C') \otimes \text{Hom}(C', C'') \rightarrow \text{Hom}(C, C'')$ is matrix multiplication. The differential $d : \text{Hom}(C, C') \rightarrow \text{Hom}(C, C')$ is defined by $df := (df_{ij}) + q'f - (-1)^l f q$ if $\deg f_{ij} = l$.

Notice that the DG category $\mathcal{A}^{\text{pre-tr}}$ is closed under formal shifts:

$$(\oplus_{i=1}^n C_i[r_i], q)[1] = (\oplus_{i=1}^n C_i[r_i + 1], -q). \quad (4.15)$$

Зсуви в dg-категорії

Given an object $A \in \mathcal{A}$, the object $A[r]$ is characterized (up to a DG isomorphism) by the existence of closed morphisms $f : A \rightarrow A[r]$, $g : A[r] \rightarrow A$ of degrees $-r$ and r , respectively, such that $fg = gf = 1$. Thus, in particular, every DG functor commutes with shifts.

Definition 4.6. The objects of $\mathcal{A}^{\text{pre-tr}}$ are “one-sided twisted complexes,” that is, formal expressions $(\oplus_{i=1}^n C_i[r_i], q)$, where $C_i \in \text{Ob } \mathcal{A}$, $r_i \in \mathbb{Z}$, $n \geq 0$, $q = (q_{ij})$, $q_{ij} \in \text{Hom}(C_j[r_j], C_i[r_i])$ is homogeneous of degree 1, $q_{ij} = 0$ for $i \geq j$, and $dq + q^2 = 0$. If $C, C' \in \text{Ob } \mathcal{A}^{\text{pre-tr}}$, $C = (\oplus C_j[r_j], q)$, $C' = (\oplus C'_j[r'_j], q')$, then the \mathbb{Z} -graded k -module $\text{Hom}(C, C')$ is the space of matrices $f = (f_{ij})$, $f_{ij} \in \text{Hom}(C_j[r_j], C'_i[r'_i])$, and the composition map $\text{Hom}(C, C') \otimes \text{Hom}(C', C'') \rightarrow \text{Hom}(C, C'')$ is matrix multiplication. The differential $d : \text{Hom}(C, C') \rightarrow \text{Hom}(C, C')$ is defined by $df := (df_{ij}) + q'f - (-1)^l f q$ if $\deg f_{ij} = l$.

Notice that the DG category $\mathcal{A}^{\text{pre-tr}}$ is closed under formal shifts:

$$(\oplus_{i=1}^n C_i[r_i], q)[1] = (\oplus_{i=1}^n C_i[r_i + 1], -q). \quad (4.15)$$

$$\begin{aligned} A \xrightarrow{f} B \xrightarrow{g} A, \quad fg = 1 = gf, \quad df = 0 = dg, \quad \deg f = -r, \quad \deg g = r, \\ A \xrightarrow{f'} C \xrightarrow{g'} A, \quad f'g' = 1 = g'f', \quad df' = 0 = dg', \quad \deg f' = -r, \quad \deg g' = r, \\ k = (B \xrightarrow{g} A \xrightarrow{f'} C), \quad j = (C \xrightarrow{g'} A \xrightarrow{f} B), \quad k, j \in Z^0 \mathcal{A}, \quad j = k^{-1} \Rightarrow B \cong C \end{aligned}$$

Конус в dg-категории

Definition 4.7. Let \mathcal{B} be a DG category and let $f \in \text{Hom}(A, B)$ be a closed degree-zero morphism in \mathcal{B} . An object $C \in \mathcal{B}$ is called the cone of f , denoted $\text{Cone}(f)$, if \mathcal{B} contains the object $A[1]$ and there exist degree-zero morphisms

$$A[1] \xrightarrow{i} C \xrightarrow{p} A[1], \quad B \xrightarrow{j} C \xrightarrow{s} B, \quad (4.16)$$

with the properties

$$pi = 1, \quad sj = 1, \quad si = 0, \quad pj = 0, \quad ip + js = 1, \quad (4.17)$$

and

$$d(j) = d(p) = 0, \quad d(i) = jf, \quad d(s) = -fp. \quad (4.18)$$

Конус в dg-категории

Definition 4.7. Let \mathcal{B} be a DG category and let $f \in \text{Hom}(A, B)$ be a closed degree-zero morphism in \mathcal{B} . An object $C \in \mathcal{B}$ is called the cone of f , denoted $\text{Cone}(f)$, if \mathcal{B} contains the object $A[1]$ and there exist degree-zero morphisms

$$A[1] \xrightarrow{i} C \xrightarrow{p} A[1], \quad B \xrightarrow{j} C \xrightarrow{s} B, \quad (4.16)$$

with the properties

$$pi = 1, \quad sj = 1, \quad si = 0, \quad pj = 0, \quad ip + js = 1, \quad (4.17)$$

and

$$d(j) = d(p) = 0, \quad d(i) = jf, \quad d(s) = -fp. \quad (4.18)$$

$$di = j \circ f \circ \sigma^{-1}, \quad ds = -f \circ \sigma^{-1} \circ p,$$

where

$$\sigma : A \rightarrow A[1], \quad \deg \sigma = -1, \quad \sigma^{-1} : A[1] \rightarrow A, \quad \deg \sigma^{-1} = 1.$$

Lemma 4.8. The cone of a closed degree-zero morphism is uniquely defined up to a DG isomorphism. \square

Proof. Note that the first set of conditions means that C is the direct sum of $A[1]$ and B in the corresponding graded category \mathcal{B}^{gr} . Thus for any object E in \mathcal{A} , there are isomorphisms of graded k -modules

$$\begin{aligned}\text{Hom}(E, C) &= \text{Hom}(E, A[1]) \oplus \text{Hom}(E, B), \\ \text{Hom}(C, E) &= \text{Hom}(A[1], E) \oplus \text{Hom}(B, E),\end{aligned}\tag{4.19}$$

which are given by composing with i and j (or with p and s). Then the second set of conditions determines the differentials in $\text{Hom}(E, C)$ and $\text{Hom}(C, E)$. \blacksquare

Lemma 4.8. The cone of a closed degree-zero morphism is uniquely defined up to a DG isomorphism. \square

Proof. Note that the first set of conditions means that C is the direct sum of $A[1]$ and B in the corresponding graded category \mathcal{B}^{gr} . Thus for any object E in \mathcal{A} , there are isomorphisms of graded k -modules

$$\begin{aligned}\text{Hom}(E, C) &= \text{Hom}(E, A[1]) \oplus \text{Hom}(E, B), \\ \text{Hom}(C, E) &= \text{Hom}(A[1], E) \oplus \text{Hom}(B, E),\end{aligned}\tag{4.19}$$

which are given by composing with i and j (or with p and s). Then the second set of conditions determines the differentials in $\text{Hom}(E, C)$ and $\text{Hom}(C, E)$. \blacksquare

We have a right dg-module $M : \mathcal{B}^{\text{op}} \rightarrow \text{dg}$,

$E \mapsto (\mathcal{B}(E, A[1]) \oplus \mathcal{B}(E, B), d_M)$, where d_M comes from the decomposition $g = i \circ (p \circ g) + j \circ (s \circ g) : E \rightarrow C$, namely,

$$\begin{aligned}d_M g &= j \circ f \circ \sigma^{-1} \circ (p \circ g) + i \circ d(p \circ g) + j \circ d(s \circ g), \\ d_M(p \circ g \oplus s \circ g) &= d(p \circ g) \oplus [f \circ \sigma^{-1} \circ (p \circ g) + d(s \circ g)],\end{aligned}$$

$$d_M = \begin{pmatrix} d & 0 \\ f \circ \sigma^{-1} \circ ? & d \end{pmatrix} \quad - \text{left matrix.}$$

Lemma 4.8. The cone of a closed degree-zero morphism is uniquely defined up to a DG isomorphism. \square

Proof. Note that the first set of conditions means that C is the direct sum of $A[1]$ and B in the corresponding graded category \mathcal{B}^{gr} . Thus for any object E in \mathcal{A} , there are isomorphisms of graded k -modules

$$\begin{aligned} \text{Hom}(E, C) &= \text{Hom}(E, A[1]) \oplus \text{Hom}(E, B), \\ \text{Hom}(C, E) &= \text{Hom}(A[1], E) \oplus \text{Hom}(B, E), \end{aligned} \tag{4.19}$$

which are given by composing with i and j (or with p and s). Then the second set of conditions determines the differentials in $\text{Hom}(E, C)$ and $\text{Hom}(C, E)$. \blacksquare

We have a right dg-module $M : \mathcal{B}^{\text{op}} \rightarrow \mathbf{dg}$,

$E \mapsto (\mathcal{B}(E, A[1]) \oplus \mathcal{B}(E, B), d_M)$, where d_M comes from the decomposition $g = i \circ (p \circ g) + j \circ (s \circ g) : E \rightarrow C$, namely,

$$d_M g = j \circ f \circ \sigma^{-1} \circ (p \circ g) + i \circ d(p \circ g) + j \circ d(s \circ g),$$

$$d_M(p \circ g \oplus s \circ g) = d(p \circ g) \oplus [f \circ \sigma^{-1} \circ (p \circ g) + d(s \circ g)],$$

$${}_E(\mathcal{E}) \quad d_M = \begin{pmatrix} d & 0 \\ f \circ \sigma^{-1} \circ ? & d \end{pmatrix} \quad - \text{left matrix.}$$

$M \cong \mathcal{B}(E, C)$ is representable (by C) iff $\exists \text{Cone}f \in \mathcal{B} (= C)$.

dg-вкладення Йонеда

LEMMA 7.3.5 (\mathcal{V} -Yoneda lemma). Given a small \mathcal{V} -category $\underline{\mathcal{D}}$, and object $d \in \underline{\mathcal{D}}$, and a \mathcal{V} -functor $F: \underline{\mathcal{D}} \rightarrow \underline{\mathcal{V}}$, the canonical map is a \mathcal{V} -natural isomorphism

$$Fd \xrightarrow{\cong} \underline{\mathcal{V}}^{\underline{\mathcal{D}}}(\underline{\mathcal{D}}(d, -), F) \longrightarrow \underline{\mathcal{V}}^{\underline{\mathcal{D}}}(\mathbb{1}, Fd)$$

\uparrow
 $\mathbb{1}$

\downarrow
 R_d

dg-вкладення Йонеда

LEMMA 7.3.5 (\mathcal{V} -Yoneda lemma). Given a small \mathcal{V} -category $\underline{\mathcal{D}}$, and object $d \in \underline{\mathcal{D}}$, and a \mathcal{V} -functor $F: \underline{\mathcal{D}} \rightarrow \underline{\mathcal{V}}$, the canonical map is a \mathcal{V} -natural isomorphism

$$Fd \xrightarrow{\cong} \underline{\mathcal{V}}^{\underline{\mathcal{D}}}(\underline{\mathcal{D}}(d, -), F).$$

Corollary

For every dg category \mathcal{C} the map defined on objects by $A \mapsto \mathcal{C}(-, A)$ extends to a fully faithful dg functor $Y_{\text{dg}}^{\mathcal{C}}: \mathcal{C} \rightarrow \text{dgMod}(\mathcal{C})$ (the dg Yoneda embedding). ~~It is easy to see that the image of~~
 $\mathcal{C}(X, Y) \rightarrow \text{dgMod}(\mathcal{C})(\mathcal{C}(_, X), \mathcal{C}(_, Y)), f \mapsto - \cdot f.$

dg-вкладення Йонеда

LEMMA 7.3.5 (\mathcal{V} -Yoneda lemma). Given a small \mathcal{V} -category $\underline{\mathcal{D}}$, and object $d \in \underline{\mathcal{D}}$, and a \mathcal{V} -functor $F: \underline{\mathcal{D}} \rightarrow \underline{\mathcal{V}}$, the canonical map is a \mathcal{V} -natural isomorphism

$$Fd \xrightarrow{\cong} \underline{\mathcal{V}}^{\underline{\mathcal{D}}}(\underline{\mathcal{D}}(d, -), F).$$

Corollary

For every dg category \mathcal{C} the map defined on objects by $A \mapsto \mathcal{C}(-, A)$ extends to a fully faithful dg functor $\mathcal{Y}_{\text{dg}}^{\mathcal{C}}: \mathcal{C} \rightarrow \text{dgMod}(\mathcal{C})$ (the dg Yoneda embedding). ~~It is easy to see that the image of~~
 $\mathcal{C}(X, Y) \rightarrow \text{dgMod}(\mathcal{C})(\mathcal{C}(_, X), \mathcal{C}(_, Y)), f \mapsto - \cdot f$.

We now give a definition of representable functor in the present situation. Let E be an object in the DG -category \mathcal{A} . It determines a contravariant DG -functor $h_E: \mathcal{A} \rightarrow C(\mathcal{A}b)$ that takes $F \in \text{Ob}\mathcal{A}$ into the complex $\text{Hom}_{\mathcal{A}}^{\bullet}(F, E)$. The assignment $E \mapsto h_E$ gives a covariant DG -functor

$$h: \mathcal{A} \rightarrow DG\text{-Fun}^0(\mathcal{A}, C(\mathcal{A}b)).$$

As in the “classical” case (see [18]), one verifies that the functor h is fully strict, i.e., that there exist isomorphisms of complexes

$$\text{Hom}_{\mathcal{A}}^{\bullet}(E, E') \simeq \text{Hom}_{DG\text{-Fun}^0(\mathcal{A}, C(\mathcal{A}b))}(h_E, h_{E'}). \quad (1.3)$$

A contravariant DG -functor $h: \mathcal{A} \rightarrow C(\mathcal{A}b)$ will be called *representable* if it is isomorphic (as a DG -functor) to a functor of the form h_E for some $E \in \text{Ob}\mathcal{A}$.

Вкладення $\text{Pre-Tr}(\mathcal{A})$ в $\text{dgMod}(\mathcal{A})$

DEFINITION 3. Let \mathcal{A} be a DG-category. We define an imbedding of DG-categories

$$\alpha: \text{Pre-Tr}(\mathcal{A}) \rightarrow \underline{\text{DG-Fun}}^0(\mathcal{A}, C(\mathcal{A}b)).$$

The imbedding assigns to an object $K = \{E_i, q_{ij}\} \in \text{Ob Pre-Tr}(\mathcal{A})$ the following DG-functor $\alpha(K): \mathcal{A} \rightarrow C(\mathcal{A}b)$. For each $E \in \text{Ob } \mathcal{A}$ the value $\alpha(K)(E)$ is the graded abelian group $\bigoplus \text{Hom}_{\mathcal{A}}(E, E_i)[i]$ provided with the differential $d+Q$, where $Q = \|q_{ij}\|$ and d is the differential in $\bigoplus \text{Hom}_{\mathcal{A}}(E, E_i)[i]$.

PROPOSITION 3. (a) *The functor α is an imbedding of $\text{Pre-Tr}(\mathcal{A})$ into $\text{DG-Fun}^0(\mathcal{A}, C(\mathcal{A}b))$ as a full DG-subcategory, and it takes the cone of a closed morphism f in $\text{Pre-Tr}(\mathcal{A})$ into the cone of the morphism $\alpha(f)$ in $\text{DG-Fun}^0(\mathcal{A}, C(\mathcal{A}b))$.*

Вкладення $\text{Pre-Tr}(\mathcal{A})$ в $\text{dgMod}(\mathcal{A})$

DEFINITION 3. Let \mathcal{A} be a DG -category. We define an imbedding of DG -categories

$$\alpha: \text{Pre-Tr}(\mathcal{A}) \rightarrow DG\text{-Fun}^0(\mathcal{A}, C(\mathcal{A}b)).$$

The imbedding assigns to an object $K = \{E_i, q_{ij}\} \in \text{Ob Pre-Tr}(\mathcal{A})$ the following DG -functor $\alpha(K): \mathcal{A} \rightarrow C(\mathcal{A}b)$. For each $E \in \text{Ob } \mathcal{A}$ the value $\alpha(K)(E)$ is the graded abelian group $\bigoplus \text{Hom}_{\mathcal{A}}(E, E_i)[i]$ provided with the differential $d+Q$, where $Q = \|q_{ij}\|$ and d is the differential in $\bigoplus \text{Hom}_{\mathcal{A}}(E, E_i)[i]$.

PROPOSITION 3. (a) *The functor α is an imbedding of $\text{Pre-Tr}(\mathcal{A})$ into $DG\text{-Fun}^0(\mathcal{A}, C(\mathcal{A}b))$ as a full DG -subcategory, and it takes the cone of a closed morphism f in $\text{Pre-Tr}(\mathcal{A})$ into the cone of the morphism $\alpha(f)$ in $DG\text{-Fun}^0(\mathcal{A}, C(\mathcal{A}b))$.*

Доведення.

Morphisms in $\text{Pre-Tr}(\mathcal{A})$ are given by rectangular matrix with entries in $\mathcal{A}(E_i, E'_j)[n'_j - n_i]$. Morphisms in $\text{dgMod}(\mathcal{A})$ are given by rectangular matrix of the same size with entries in $\text{dgMod}(\mathcal{A})(\mathcal{A}(_, E_i)[n_i], \mathcal{A}(_, E'_j)[n'_j])$. Differentials agree. \square

Трикутники в dg-категорії з конусами

Given a closed degree-zero morphism $f : A \rightarrow B$, the diagram

$$A \xrightarrow{f} B \xrightarrow{j} \text{Cone}(f) \xrightarrow{p} A[1] \quad (4.20)$$

is called a pre-exact triangle.

Remark 4.9. It is clear that any DG functor preserves cones of closed degree-zero morphisms and preserves pre-exact triangles.

Замкненість $\mathcal{A}^{\text{pre-tr}}$ стосовно конусів

Proposition 4.10 [5]. Let \mathcal{A} be a DG category. Then

- (a) the DG category $\mathcal{A}^{\text{pre-tr}}$ is closed under taking cones of closed degree-zero morphisms;
- (b) every object in $\mathcal{A}^{\text{pre-tr}}$ can be obtained from objects in \mathcal{A} by taking successive cones of closed degree-zero morphisms. \square

Proof. (a) Given a closed morphism of degree zero

$$f : (\oplus C_i[r_i], q) \longrightarrow (\oplus C'_j[r'_j], q'), \quad (4.21)$$

its cone is the twisted complex $(\oplus C'_j[r'_j] \oplus C_i[r_i + 1], (q', -q + f))$. For example, if $A, B \in \mathcal{A}$ and $f : A \rightarrow B$ is a closed morphism of degree zero, then $\text{Cone}(f)$ is the twisted complex $(B \oplus A[1], (0, f)) \in \mathcal{A}^{\text{pre-tr}}$.

(b) Let $C = (\oplus_{i=1}^n C_i[r_i], q)$ be a twisted complex. Consider its twisted subcomplex $C' = (\oplus_{i=1}^{n-1} C_i[r_i], q')$, where $q' = q - \oplus_i q_{in}$. Then C is the cone of the closed degree-zero morphism $\oplus_{i=1}^{n-1} q_{in} : (C_n[r_n - 1], 0) \rightarrow C'$. \blacksquare

Замкненість $\mathcal{A}^{\text{pre-tr}}$ стосовно конусів

Proposition 4.10 [5]. Let \mathcal{A} be a DG category. Then

- (a) the DG category $\mathcal{A}^{\text{pre-tr}}$ is closed under taking cones of closed degree-zero morphisms;
- (b) every object in $\mathcal{A}^{\text{pre-tr}}$ can be obtained from objects in \mathcal{A} by taking successive cones of closed degree-zero morphisms. \square

Proof. (a) Given a closed morphism of degree zero

$$f : (\oplus C_i[r_i], q) \longrightarrow (\oplus C'_j[r'_j], q'), \quad (4.21)$$

its cone is the twisted complex $(\oplus C'_j[r'_j] \oplus C_i[r_i + 1], (q', -q + f))$. For example, if $A, B \in \mathcal{A}$ and $f : A \rightarrow B$ is a closed morphism of degree zero, then $\text{Cone}(f)$ is the twisted complex $(B \oplus A[1], (0, f)) \in \mathcal{A}^{\text{pre-tr}}$.

(b) Let $C = (\oplus_{i=1}^n C_i[r_i], q)$ be a twisted complex. Consider its twisted subcomplex $C' = (\oplus_{i=1}^{n-1} C_i[r_i], q')$, where $q' = q - \oplus_i q_{i,n}$. Then C is the cone of the closed degree-zero morphism $\oplus_{i=1}^{n-1} q_{i,n} : (C_n[r_n - 1], 0) \rightarrow C'$. \blacksquare

$$\left(\begin{array}{c|c} q' & 0 \\ \hline f \circ \sigma^{-1} & -q \end{array} \right)$$

(Сильно) передтриангульовані dg-категорії

A DG category \mathcal{A} is said to be *pretriangulated* if for every $A \in \mathcal{A}$, $k \in \mathbb{Z}$, the object $A[k] \in \mathcal{A}^{\text{pre-tr}}$ is homotopy-equivalent to an object of \mathcal{A} and for every closed morphism of degree-zero f in \mathcal{A} , the object $\text{Cone}(f) \in \mathcal{A}^{\text{pre-tr}}$ is homotopy-equivalent to an object of \mathcal{A} . We say that \mathcal{A} is *strongly pretriangulated* if the same is true with “homotopy-equivalent” replaced by “DG isomorphic.” Actually, if \mathcal{A} is pretriangulated (resp., strongly pretriangulated), then every object of $\mathcal{A}^{\text{pre-tr}}$ is homotopy-equivalent (resp., DG isomorphic) to an object of \mathcal{A} [9]. Thus, \mathcal{A} is pretriangulated (resp., strongly pretriangulated) if and only if the embedding $\text{Ho}(\mathcal{A}) \hookrightarrow \text{Ho}(\mathcal{A}^{\text{pre-tr}})$ is an equivalence (resp., the embedding $\mathcal{A} \hookrightarrow \mathcal{A}^{\text{pre-tr}}$ is a DG equivalence).

(Сильно) передтриангульовані dg-категорії

A DG category \mathcal{A} is said to be *pretriangulated* if for every $A \in \mathcal{A}$, $k \in \mathbb{Z}$, the object $A[k] \in \mathcal{A}^{\text{pre-tr}}$ is homotopy-equivalent to an object of \mathcal{A} and for every closed morphism of degree-zero f in \mathcal{A} , the object $\text{Cone}(f) \in \mathcal{A}^{\text{pre-tr}}$ is homotopy-equivalent to an object of \mathcal{A} . We say that \mathcal{A} is *strongly pretriangulated* if the same is true with “homotopy-equivalent” replaced by “DG isomorphic.” Actually, if \mathcal{A} is pretriangulated (resp., strongly pretriangulated), then every object of $\mathcal{A}^{\text{pre-tr}}$ is homotopy-equivalent (resp., DG isomorphic) to an object of \mathcal{A} [9]. Thus, \mathcal{A} is pretriangulated (resp., strongly pretriangulated) if and only if the embedding $\text{Ho}(\mathcal{A}) \hookrightarrow \text{Ho}(\mathcal{A}^{\text{pre-tr}})$ is an equivalence (resp., the embedding $\mathcal{A} \hookrightarrow \mathcal{A}^{\text{pre-tr}}$ is a DG equivalence).

Definition 3.3. A dg category \mathcal{C} is *strongly pretriangulated* if $A[n]$ and $\text{Cone}(f)$ exist (in \mathcal{C}), for every $n \in \mathbb{Z}$, every object A of \mathcal{C} and every morphism f of $Z^0(\mathcal{C})$.

A dg category \mathcal{C} is *pretriangulated* if there exists a quasi-equivalence $\mathcal{C} \rightarrow \mathcal{C}'$ with \mathcal{C}' strongly pretriangulated.

$\text{dgMod}(\mathcal{C})$ – сильно передтриангульована dg-категорія

Let $\phi : M \rightarrow N \in Z^0 \text{dgMod}(\mathcal{C})$. It is given by the family $\phi(X) : M(X) \rightarrow N(X) \in \mathbf{dg}$ such that for all $f \in \mathcal{C}(X, Y)$

$$\begin{array}{ccc} M(X) & \xrightarrow{M(f)} & M(Y) \\ \phi(X) \downarrow & = & \downarrow \phi(Y) \\ N(X) & \xrightarrow{N(f)} & N(Y) \end{array}$$

$\text{dgMod}(\mathcal{C})$ – сильно передтриангульована dg-категорія

Let $\phi : M \rightarrow N \in Z^0 \text{dgMod}(\mathcal{C})$. It is given by the family $\phi(X) : M(X) \rightarrow N(X) \in \mathbf{dg}$ such that for all $f \in \mathcal{C}(X, Y)$

$$\begin{array}{ccc} M(X) & \xrightarrow{M(f)} & M(Y) \\ \phi(X) \downarrow & = & \downarrow \phi(Y) \\ N(X) & \xrightarrow{N(f)} & N(Y) \end{array}$$

Define \mathcal{C} -module $\mathbf{Cone} \phi$ by

$(\mathbf{Cone} \phi)(X) = \mathbf{Cone}(\phi(X)) \equiv \mathbf{Cone} \phi(X) = (M(X)[1] \oplus N(X), d)$
and $\mathcal{C}^{\text{op}}(X, Y) \rightarrow \underline{\mathbf{dg}}(\mathbf{Cone} \phi(X), \mathbf{Cone} \phi(Y)), f \mapsto M(f)[1] \oplus N(f)$.

$\text{dgMod}(\mathcal{C})$ – сильно передтриангульована dg-категорія

Let $\phi : M \rightarrow N \in Z^0 \text{dgMod}(\mathcal{C})$. It is given by the family $\phi(X) : M(X) \rightarrow N(X) \in \mathbf{dg}$ such that for all $f \in \mathcal{C}(X, Y)$

$$\begin{array}{ccc} M(X) & \xrightarrow{M(f)} & M(Y) \\ \phi(X) \downarrow & = & \downarrow \phi(Y) \\ N(X) & \xrightarrow{N(f)} & N(Y) \end{array}$$

Define \mathcal{C} -module $\mathbf{Cone} \phi$ by

$(\mathbf{Cone} \phi)(X) = \mathbf{Cone}(\phi(X)) \equiv \mathbf{Cone} \phi(X) = (M(X)[1] \oplus N(X), d)$
and $\mathcal{C}^{\text{op}}(X, Y) \rightarrow \underline{\mathbf{dg}}(\mathbf{Cone} \phi(X), \mathbf{Cone} \phi(Y)), f \mapsto M(f)[1] \oplus N(f)$.

Exercise

Verify that this is a chain map.

Ортогональні категорії

6-2 Proposition : Soit N une sous-catégorie triangulée d'une catégorie triangulée A . La catégorie pleine N^\perp (resp ${}^\perp N$) engendrée par les objets X de A tels que pour tout objet Y de N on ait $\text{Hom}_A(Y,X) = 0$ (resp $\text{Hom}_A(X,Y) = 0$), est une sous-catégorie épaisse de A . La catégorie N^\perp est appelée par abus de langage, l'orthogonale à droite de N .

Ортогональні категорії

6-2 Proposition : Soit N une sous-catégorie triangulée d'une catégorie triangulée A . La catégorie pleine N^\perp (resp ${}^\perp N$) engendrée par les objets X de A tels que pour tout objet Y de N on ait $\text{Hom}_A(Y,X) = 0$ (resp $\text{Hom}_A(X,Y) = 0$), est une sous-catégorie épaisse de A . La catégorie N^\perp est appelée par abus de langage, l'orthogonale à droite de N .

1-1 Définition : Une sous-catégorie B d'une catégorie triangulée A est dite épaisse si B est une sous-catégorie triangulée pleine de A et si de plus B possède la propriété suivante :

Pour tout morphisme $f : X \rightarrow Y$, se factorisant par un objet de B et contenu dans un triangle distingué (X,Y,Z,f,g,h) où Z est un objet de B , la source de f et le but de f sont des objets de B .

Ортогональні категорії

6-2 Proposition : Soit N une sous-catégorie triangulée d'une catégorie triangulée A . La catégorie pleine N^\perp (resp ${}^\perp N$) engendrée par les objets X de A tels que pour tout objet Y de N on ait $\text{Hom}_A(Y, X) = 0$ (resp $\text{Hom}_A(X, Y) = 0$), est une sous-catégorie épaisse de A . La catégorie N^\perp est appelée par abus de langage, l'orthogonale à droite de N .

1-1 Définition : Une sous-catégorie B d'une catégorie triangulée A est dite épaisse si B est une sous-catégorie triangulée pleine de A et si de plus B possède la propriété suivante :

Pour tout morphisme $f : X \rightarrow Y$, se factorisant par un objet de B et contenu dans un triangle distingué (X, Y, Z, f, g, h) où Z est un objet de B , la source de f et le but de f sont des objets de B .

(ii) *For any $W \in \text{Ob}(\mathcal{C})$, $\text{Hom}_{\mathcal{C}}(W, \cdot)$ and $\text{Hom}_{\mathcal{C}}(\cdot, W)$ are cohomological functors.*

Proof of Proposition 6-2.

Let $\mathcal{N} \subset \mathcal{A}$ be a full subcategory closed wrt shifts. Let a full subcategory $\mathcal{L} = {}^{\perp}\mathcal{N}$ consist of $X \in \mathbf{Ob} \mathcal{A}$ s/t $\mathcal{A}(X, \mathcal{N}) = 0$.

$\forall N \in \mathcal{N}$ and for any distinguished triangle $X \rightarrow Y \rightarrow Z \rightarrow$ in \mathcal{A} there is an exact sequence

$$\mathcal{A}(X, N[-1]) \rightarrow \mathcal{A}(Z, N) \rightarrow \mathcal{A}(Y, N) \rightarrow \mathcal{A}(X, N) \rightarrow \mathcal{A}(Z, N[1]).$$

Proof of Proposition 6-2.

Let $\mathcal{N} \subset \mathcal{A}$ be a full subcategory closed wrt shifts. Let a full subcategory $\mathcal{L} = {}^{\perp}\mathcal{N}$ consist of $X \in \mathbf{Ob} \mathcal{A}$ s/t $\mathcal{A}(X, \mathcal{N}) = 0$.

$\forall N \in \mathcal{N}$ and for any distinguished triangle $X \rightarrow Y \rightarrow Z \rightarrow$ in \mathcal{A} there is an exact sequence

$$\mathcal{A}(X, N[-1]) \rightarrow \mathcal{A}(Z, N) \rightarrow \mathcal{A}(Y, N) \rightarrow \mathcal{A}(X, N) \rightarrow \mathcal{A}(Z, N[1]).$$

If two of three vertices of the triangle are in \mathcal{L} so is the third.
Thus \mathcal{L} is a triangulated subcategory of \mathcal{A} .

Proof of Proposition 6-2.

Let $\mathcal{N} \subset \mathcal{A}$ be a full subcategory closed wrt shifts. Let a full subcategory $\mathcal{L} = {}^{\perp}\mathcal{N}$ consist of $X \in \mathbf{Ob} \mathcal{A}$ s/t $\mathcal{A}(X, \mathcal{N}) = 0$.

$\forall N \in \mathcal{N}$ and for any distinguished triangle $X \rightarrow Y \rightarrow Z \rightarrow$ in \mathcal{A} there is an exact sequence

$$\mathcal{A}(X, N[-1]) \rightarrow \mathcal{A}(Z, N) \rightarrow \mathcal{A}(Y, N) \rightarrow \mathcal{A}(X, N) \rightarrow \mathcal{A}(Z, N[1]).$$

If two of three vertices of the triangle are in \mathcal{L} so is the third. Thus \mathcal{L} is a triangulated subcategory of \mathcal{A} .

Assume that $Z \in \mathcal{L}$, $N \in \mathcal{N}$ and $f : X \rightarrow Y$ factorises through $L \in \mathcal{L}$. Then the exact sequence implies that

$\mathcal{A}(f, N) : \mathcal{A}(Y, N) \rightarrow \mathcal{A}(X, N)$ is a bijection. On the other hand,

$$\mathcal{A}(f, N) = (\mathcal{A}(Y, N) \rightarrow \mathcal{A}(L, N) \rightarrow \mathcal{A}(X, N)) = 0.$$

Proof of Proposition 6-2.

Let $\mathcal{N} \subset \mathcal{A}$ be a full subcategory closed wrt shifts. Let a full subcategory $\mathcal{L} = {}^{\perp}\mathcal{N}$ consist of $X \in \mathbf{Ob} \mathcal{A}$ s/t $\mathcal{A}(X, \mathcal{N}) = 0$.

$\forall N \in \mathcal{N}$ and for any distinguished triangle $X \rightarrow Y \rightarrow Z \rightarrow$ in \mathcal{A} there is an exact sequence

$$\mathcal{A}(X, N[-1]) \rightarrow \mathcal{A}(Z, N) \rightarrow \mathcal{A}(Y, N) \rightarrow \mathcal{A}(X, N) \rightarrow \mathcal{A}(Z, N[1]).$$

If two of three vertices of the triangle are in \mathcal{L} so is the third. Thus \mathcal{L} is a triangulated subcategory of \mathcal{A} .

Assume that $Z \in \mathcal{L}$, $N \in \mathcal{N}$ and $f : X \rightarrow Y$ factorises through $L \in \mathcal{L}$. Then the exact sequence implies that $\mathcal{A}(f, N) : \mathcal{A}(Y, N) \rightarrow \mathcal{A}(X, N)$ is a bijection. On the other hand,

$$\mathcal{A}(f, N) = (\mathcal{A}(Y, N) \rightarrow \mathcal{A}(L, N) \rightarrow \mathcal{A}(X, N)) = 0.$$

Hence, $\mathcal{A}(X, N) = 0$ and $\mathcal{A}(Y, N) = 0$. Therefore, \mathcal{L} is épaisse. □

Досконалі модулі

If \mathcal{C} is a dg category, $\text{dgMod}(\mathcal{C})$, $\text{dgAcy}(\mathcal{C})$ and $\text{h-proj}(\mathcal{C})$ are strongly pretriangulated dg categories. Moreover, the (triangulated) categories $H^0(\text{dgMod}(\mathcal{C}))$, $H^0(\text{dgAcy}(\mathcal{C}))$ and $H^0(\text{h-proj}(\mathcal{C}))$ have arbitrary coproducts, and there is a semi-orthogonal decomposition

$$(3.1) \quad H^0(\text{dgMod}(\mathcal{C})) = \langle H^0(\text{dgAcy}(\mathcal{C})), H^0(\text{h-proj}(\mathcal{C})) \rangle.$$

This clearly implies that there is an exact equivalence between $H^0(\text{h-proj}(\mathcal{C}))$ and the Verdier quotient $\mathcal{D}(\mathcal{C}) := H^0(\text{dgMod}(\mathcal{C}))/H^0(\text{dgAcy}(\mathcal{C}))$ (which is by definition the *derived category* of \mathcal{C}).

For every dg category \mathcal{C} we will denote by $\text{Pretr}(\mathcal{C})$ (respectively, $\text{Perf}(\mathcal{C})$) the smallest full dg subcategory of $\text{h-proj}(\mathcal{C})$ containing $\mathbf{Y}_{\text{dg}}^{\mathcal{C}}(\mathcal{C})$ and closed under homotopy equivalences, shifts, cones (respectively, also direct summands in $H^0(\text{h-proj}(\mathcal{C}))$). It is easy to see that $\text{Pretr}(\mathcal{C})$ and $\text{Perf}(\mathcal{C})$ are strongly pretriangulated and that \mathcal{C} is pretriangulated if and only if $\mathbf{Y}_{\text{dg}}^{\mathcal{C}} : \mathcal{C} \rightarrow \text{Pretr}(\mathcal{C})$ is a quasi-equivalence. Moreover, $\text{Pretr}(\mathcal{C}) \subseteq \text{Perf}(\mathcal{C})$ and $H^0(\text{Perf}(\mathcal{C}))$ can be identified with the idempotent completion $H^0(\text{Pretr}(\mathcal{C}))^{\text{ic}}$ of $H^0(\text{Pretr}(\mathcal{C}))$. Hence $\mathbf{Y}_{\text{dg}}^{\mathcal{C}} : \mathcal{C} \rightarrow \text{Perf}(\mathcal{C})$ is a quasi-equivalence if and only if \mathcal{C} is pretriangulated and $H^0(\mathcal{C})$ is idempotent complete.

Remark 3.5. Recall that an additive category \mathcal{A} is *idempotent complete* if every idempotent (namely, a morphism $e : A \rightarrow A$ in \mathcal{A} such that $e^2 = e$) splits, or, equivalently, has a kernel. Every additive category \mathcal{A} admits a fully faithful and additive embedding $\mathcal{A} \hookrightarrow \mathcal{A}^{\text{ic}}$, where \mathcal{A}^{ic} is an idempotent complete additive category, with the property that every object of \mathcal{A}^{ic} is a direct summand of an object from \mathcal{A} . The category \mathcal{A}^{ic} (or, better, the functor $\mathcal{A} \rightarrow \mathcal{A}^{\text{ic}}$) is called the *idempotent completion* of \mathcal{A} . It can be proved (see [2]) that, if \mathcal{T} is a triangulated category, then \mathcal{T}^{ic} is triangulated as well (and $\mathcal{T} \hookrightarrow \mathcal{T}^{\text{ic}}$ is exact).

Досконалі модулі

If \mathcal{C} is a dg category, $\mathrm{dgMod}(\mathcal{C})$, $\mathrm{dgAcy}(\mathcal{C})$ and $\mathrm{h-proj}(\mathcal{C})$ are strongly pretriangulated dg categories. Moreover, the (triangulated) categories $H^0(\mathrm{dgMod}(\mathcal{C}))$, $H^0(\mathrm{dgAcy}(\mathcal{C}))$ and $H^0(\mathrm{h-proj}(\mathcal{C}))$ have arbitrary coproducts, and there is a semi-orthogonal decomposition

$$(3.1) \quad H^0(\mathrm{dgMod}(\mathcal{C})) = \langle H^0(\mathrm{dgAcy}(\mathcal{C})), H^0(\mathrm{h-proj}(\mathcal{C})) \rangle.$$

This clearly implies that there is an exact equivalence between $H^0(\mathrm{h-proj}(\mathcal{C}))$ and the Verdier quotient $\mathcal{D}(\mathcal{C}) := H^0(\mathrm{dgMod}(\mathcal{C}))/H^0(\mathrm{dgAcy}(\mathcal{C}))$ (which is by definition the *derived category* of \mathcal{C}).

For every dg category \mathcal{C} we will denote by $\mathrm{Pretr}(\mathcal{C})$ (respectively, $\mathrm{Perf}(\mathcal{C})$) the smallest full dg subcategory of $\mathrm{h-proj}(\mathcal{C})$ containing $\mathsf{Y}_{\mathrm{dg}}^{\mathcal{C}}(\mathcal{C})$ and closed under homotopy equivalences, shifts, cones (respectively, also direct summands in $H^0(\mathrm{h-proj}(\mathcal{C}))$). It is easy to see that $\mathrm{Pretr}(\mathcal{C})$ and $\mathrm{Perf}(\mathcal{C})$ are strongly pretriangulated and that \mathcal{C} is pretriangulated if and only if $\mathsf{Y}_{\mathrm{dg}}^{\mathcal{C}} : \mathcal{C} \rightarrow \mathrm{Pretr}(\mathcal{C})$ is a quasi-equivalence. Moreover, $\mathrm{Pretr}(\mathcal{C}) \subseteq \mathrm{Perf}(\mathcal{C})$ and $H^0(\mathrm{Perf}(\mathcal{C}))$ can be identified with the idempotent completion $H^0(\mathrm{Pretr}(\mathcal{C}))^{\mathrm{ic}}$ of $H^0(\mathrm{Pretr}(\mathcal{C}))$. Hence $\mathsf{Y}_{\mathrm{dg}}^{\mathcal{C}} : \mathcal{C} \rightarrow \mathrm{Perf}(\mathcal{C})$ is a quasi-equivalence if and only if \mathcal{C} is pretriangulated and $H^0(\mathcal{C})$ is idempotent complete.

Remark 3.5. Recall that an additive category \mathcal{A} is *idempotent complete* if every idempotent (namely, a morphism $e : A \rightarrow A$ in \mathcal{A} such that $e^2 = e$) splits, or, equivalently, has a kernel. Every additive category \mathcal{A} admits a fully faithful and additive embedding $\mathcal{A} \hookrightarrow \mathcal{A}^{\mathrm{ic}}$, where $\mathcal{A}^{\mathrm{ic}}$ is an idempotent complete additive category, with the property that every object of $\mathcal{A}^{\mathrm{ic}}$ is a direct summand of an object from \mathcal{A} . The category $\mathcal{A}^{\mathrm{ic}}$ (or, better, the functor $\mathcal{A} \rightarrow \mathcal{A}^{\mathrm{ic}}$) is called the *idempotent completion* of \mathcal{A} . It can be proved (see [2]) that, if \mathcal{T} is a triangulated category, then $\mathcal{T}^{\mathrm{ic}}$ is triangulated as well (and $\mathcal{T} \hookrightarrow \mathcal{T}^{\mathrm{ic}}$ is exact).

$$\forall N \in \mathrm{dgMod}(\mathcal{C}) \exists M \in \mathrm{h-proj}(\mathcal{C}) \exists M \xrightarrow{f} N \rightarrow \mathrm{Cone} f, \in \mathrm{dgAcy}(\mathcal{C})$$

Досконалі комплекси

A **perfect complex** of modules over a commutative ring \mathbb{k} is an object in the derived category of \mathbb{k} -modules that is quasi-isomorphic to a bounded complex of finitely generated projective \mathbb{k} -modules.

Досконалі комплекси

A **perfect complex** of modules over a commutative ring \mathbb{k} is an object in the derived category of \mathbb{k} -modules that is quasi-isomorphic to a bounded complex of finitely generated projective \mathbb{k} -modules.

Perfect complexes are precisely the compact objects in the unbounded derived category $D(\mathbb{k})$ of \mathbb{k} -modules.

Досконалі комплекси

A **perfect complex** of modules over a commutative ring \mathbb{k} is an object in the derived category of \mathbb{k} -modules that is quasi-isomorphic to a bounded complex of finitely generated projective \mathbb{k} -modules.

Perfect complexes are precisely the compact objects in the unbounded derived category $D(\mathbb{k})$ of \mathbb{k} -modules.

An object X in a category \mathcal{C} which admits all filtered colimits is called **compact** if the functor

$$\mathcal{C}(X, \cdot) : \mathcal{C} \rightarrow \text{Sets}, Y \mapsto \mathcal{C}(X, Y)$$

commutes with filtered colimits, i.e., if the natural map

$$\text{colim } \mathcal{C}(X, Y_i) \rightarrow \mathcal{C}(X, \text{colim}_i Y_i)$$

is a bijection for any filtered system of objects Y_i in \mathcal{C} .

Досконалі комплекси

A **perfect complex** of modules over a commutative ring \mathbb{k} is an object in the derived category of \mathbb{k} -modules that is quasi-isomorphic to a bounded complex of finitely generated projective \mathbb{k} -modules.

Perfect complexes are precisely the compact objects in the unbounded derived category $D(\mathbb{k})$ of \mathbb{k} -modules.

An object X in a category \mathcal{C} which admits all filtered colimits is called **compact** if the functor

$$\mathcal{C}(X, \cdot) : \mathcal{C} \rightarrow \text{Sets}, Y \mapsto \mathcal{C}(X, Y)$$

commutes with filtered colimits, i.e., if the natural map

$$\text{colim } \mathcal{C}(X, Y_i) \rightarrow \mathcal{C}(X, \text{colim}_i Y_i)$$

is a bijection for any filtered system of objects Y_i in \mathcal{C} .

Since elements in the filtered colimit at the left are represented by maps $X \rightarrow Y_i$, for some i , the surjectivity of the above map amounts to requiring that a map $X \rightarrow \text{colim}_i Y_i$ factors over some Y_i .

Добре породжені триангульовані категорії

Let \mathcal{T} be a triangulated category with small coproducts. For a cardinal α , an object S of \mathcal{T} is α -small if every map $S \rightarrow \coprod_{i \in I} X_i$ in \mathcal{T} (where I is a small set) factors through $\coprod_{i \in J} X_i$, for some $J \subseteq I$ with $|J| < \alpha$. A cardinal α is called regular if it is not the sum of fewer than α cardinals, all of them smaller than α .





Добре породжені триангульовані категорії




Let \mathcal{T} be a triangulated category with small coproducts. For a cardinal α , an object S of \mathcal{T} is α -small if every map $S \rightarrow \coprod_{i \in I} X_i$ in \mathcal{T} (where I is a small set) factors through $\coprod_{i \in J} X_i$, for some $J \subseteq I$ with $|J| < \alpha$. A cardinal α is called regular if it is not the sum of fewer than α cardinals, all of them smaller than α .

Definition

The category \mathcal{T} is well generated if there exists a small set \mathcal{S} of objects in \mathcal{T} satisfying the following properties:

- (G1) An object $X \in \mathcal{T}$ is isomorphic to 0, if and only if $\mathcal{T}(S, X[j]) = 0$, for all $S \in \mathcal{S}$ and all $j \in \mathbb{Z}$;
- (G2) For every small set of maps $\{X_i \rightarrow Y_i\}_{i \in I}$ in \mathcal{T} , the induced map $\mathcal{T}(S, \coprod_i X_i) \rightarrow \mathcal{T}(S, \coprod_i Y_i)$ is surjective for all $S \in \mathcal{S}$, if $\mathcal{T}(S, X_i) \rightarrow \mathcal{T}(S, Y_i)$ is surjective, for all $i \in I$ and all $S \in \mathcal{S}$;
- (G3) There exists a regular cardinal α such that every object of \mathcal{S} is α -small.

-  Alexei I. Bondal and Mikhail M. Kapranov, Enhanced triangulated categories, *Mat. Sb.* 181 (1990), no. 5, 669–683, English transl. in *Math. USSR Sbornik*, 70 (1991), no. 1, 93–107. Definition 1.3
-  Alexei I. Bondal, Michael Larsen, and Valery A. Lunts, Grothendieck ring of pretriangulated categories, *Int. Math. Res. Not.* (2004), no. 29, 1461–1495, arXiv:math.AG/0401009. Definition 4.6 – Proposition 4.10
-  Alberto Canonaco, Amnon Neeman, and Paolo Stellari, Uniqueness of enhancements for derived and geometric categories, 2021, arXiv:2101.04404 §3.1 – §3.5
-  Masaki Kashiwara and Pierre Schapira, Sheaves on manifolds, *Grundlehren der mathematischen Wissenschaften*, vol. 292, Springer-Verlag, Berlin, New York, 1990. Proposition 1.5.3(ii)

-  Emily Riehl, *Categorical homotopy theory*, New Mathematical Monographs, vol. 24, Cambridge University Press, Cambridge, 2014, <http://gen.lib.rus.ec/book/index.php?md5=d20d520ad0551d9e0cf2072b168aac1c>. Lemma 7.3.5
-  Gonçalo Tabuada, Une structure de catégorie de modèles de Quillen sur la catégorie des dg-catégories, *C. R. Math. Acad. Sci. Paris* 340 (2005), no. 1, 15–19, [arXiv:math.KT/0407338](https://arxiv.org/abs/math/0407338) Theorem 0.1
-  Jean-Louis Verdier, *Catégories dérivées: Quelques résultats (État 0)*, *Cohomologie Étale*, *Lecture Notes in Math.*, no. 569, Springer-Verlag, 1977, pp. 262–311. Proposition 1.2.6-2